

Betaine for homocystinuria

GP Information Sheet

Introduction

Betaine occurs naturally in the body as a metabolite and is present in small amounts in foods such as sugarbeet, spinach, cereals, and seafood. Betaine is a licensed drug (some unlicensed versions are also available) which has been used for many years on a named-patient basis as a supplement to other therapies such as vitamin B6, vitamin B12, folate and a specific diet for the treatment of classical homocystinuria in neonates, infants, children and adults. There are no licensed comparator products available.

Classical homocystinuria is a genetic disorder of metabolism of the amino acid methionine, used by the body for growth and repair. Methionine is normally broken down to the amino acid homocysteine and then to cysteine. Patients with homocystinuria are missing an enzyme that converts homocysteine into cysteine in the liver leading to very elevated plasma homocysteine levels. This imbalance can lead to mental retardation, ocular abnormalities, osteoporosis, premature atherosclerosis and thromboembolic disease. There are two different pathways of metabolism of homocysteine. To treat homocystinuria, excess homocysteine is remethylated to methionine, and Betaine acts as the methyl group donor in this process.

Dosage and Administration

Adults and Children >10 yrs old: 3g orally twice daily adjusted according to response; max 10g twice daily.
Children <10 yrs old: 50mg/kg twice daily, dose and frequency adjusted according to response; max 75mg/kg twice daily.

The dosage is increased to lower total homocysteine levels and reduce free homocysteine to undetectable levels. Response usually occurs within a month.

Available as

| Route | Dosage Forms | Strength | Brand Names | Supplier |
|-------|--------------------------|------------------------------------------------|-------------|-----------------------------------------|
| Oral | Powder for oral solution | 500mg/1ml when reconstituted | Unlicensed | Veriton Pharma 01932 690325 |
| Oral | Powder for oral solution | 100mg, 150mg or 1g per level scoopful provided | Cystadane® | Orphan Medical UK Tel: 01491 414 333 |

Monitoring Requirements

Monitoring is performed by the specialist at clinic appointments which are generally every 6 months. This involves:

- measurement of the plasma concentration of homocysteine (to determine the response to treatment and any dose adjustment required)
- measurement of plasma methionine levels at start of treatment and periodically thereafter (as cerebral oedema has been reported in patients with hypermethioninaemia).

Adverse Effects, Precautions and Contraindications

Increased levels of methionine are commonly reported.

The following are all listed as occurring uncommonly: anorexia, agitation, depression, irritability, dental disorders, diarrhoea, glossitis, nausea, stomach discomfort, vomiting, hair loss, abnormal skin odour, urinary incontinence. Uncommon cases of severe cerebral oedema have been reported after starting therapy with complete recovery after treatment discontinuation.

Common Drug Interactions

No interaction studies have been performed. However, based on in vitro data, betaine might interact with amino acid mixtures and medicinal products containing vigabatrin and GABA analogues.

Communication

For any queries relating to this patient's treatment with betaine, please contact the specialist.

This information is not inclusive of all prescribing information and potential adverse effects.

Please refer to full prescribing data in the SPC or the BNF
Date Prepared: March 2021 Date of review: March 2026